

## RESEARCH COMMUNICATION

# Ethnicity Greatly Influences the Interleukin-1 Gene Cluster(IL-1 $\beta$ Promoter, Exon-5 and IL-1Ra) Polymorphisms: A Pilot Study of a North Indian Population

Parmeet Kaur Manchanda, Hemant K Bid, Rama D Mittal\*

## Abstract

There is considerable evidence that polymorphisms in the regulatory regions of cytokine genes are highly influenced by ethnicity. Polymorphisms in interleukin 1- $\beta$  (IL-1 $\beta$ ) and IL-1 receptor antagonist (IL-1Ra) genes, respectively encoding a potent inflammatory agent and an antagonist, which combines with IL-1 receptors competitively, have been associated with a number of diseases like systemic lupus erythematosus, rheumatoid arthritis, sepsis, kidney diseases, and cancer. In this study, we therefore evaluated the distribution of interleukin-1 gene cluster (IL-1 $\beta$  promoter region, exon-5 and IL-1Ra) gene polymorphisms in 206 healthy north Indian subjects, using PCR-based restriction analysis. We also constructed various haplotypes and estimated the linkage disequilibrium (LD). We found that genotype and allelic frequencies for these cytokines were conspicuously different when compared among different ethnic populations. The haplotype 'T-E1-1' predominated (41.7%) while the least common was 'C-E2-2' (2%) in our population. Genetic linkage between three loci of IL-1 gene showed strong association among the variants in controls ( $D' = 0.42$ ,  $p < 0.001$ ). Our results suggest that the frequency and distribution of the polymorphisms in India are substantially different from other populations and ethnic groups. Thus they signify an impact of ethnicity and provide a basis for future epidemiological and clinical studies.

**Key Words:** IL-1 gene cluster - SNPs - PCR-RFLP - ethnicity

*Asian Pacific J Cancer Prev*, 6, 541-546

## Introduction

DNA sequences of the human genome reveal that many genes are polymorphic. In coding or noncoding regions of a specific gene, there may be either a single base pair substitution of one nucleotide for another (SNPs) or a variable number of repeats of a short repetitive DNA sequence (VNTR). Gene-environment interactions may be manifested in various ways, either by risk effects based on an individual's genotype, or differential gene risk effects based on exposure (Michael et al., 2002). The study of genetic polymorphisms promises to help define pathophysiological mechanisms, to identify individuals at risk for disease and to suggest novel targets for drug treatment.

Cytokines are signaling molecules contributing to the inflammatory response, and are key components in the pathogenesis of many diseases like cancer, metabolic disorders and inflammatory conditions (Anderson et al.,

2001). Interleukin-1 $\beta$ , located at chromosome 2q12, is a potent proinflammatory agent that is central in immunoregulation, inflammation and cancer formation (Cantagrel et al., 1999). IL-1 receptor antagonist (IL-Ra), a structural variant of IL-1, binds to the same IL-1 receptor and acts as a competitive inhibitor of IL-1 bioactivity. In the second intron of the IL-1RN gene, there is a functional VNTR polymorphism, which is characterized as having an important role in regulating the serum IL-Ra levels, human immune response and cancer risk. (Hu et al., 2005). Interleukin (IL)-1 is known to act as a tumor growth factor by inducing angiogenic factors (Konishi et al., 2005).

Genetic variation in the human genome is an emerging resource for studying cancer, a complex set of diseases characterized by both environmental and genetic contributions. Risks of cancer are clearly influenced by polymorphisms in genes involved in carcinogen metabolism and the immune system. The primary focus of this article is the problem of identifying DNA sequence variation that is

Department of Urology, Sanjay Gandhi Post Graduate Institute of Medical Sciences, Lucknow-226014 \* Correspondence R.D. Mittal, Additional Professor, Department of Urology, Sanjay Gandhi Postgraduate Institute of Medical Sciences, Lucknow-226014, India Fax: 91-522-2668017 Email: rmittal@sgpgi.ac.in, ramamittal@yahoo.com

segregated in a population and has a causal connection with complex disease like cancer. One could perform tests of this association without consideration of the fact that the SNPs are not independent of one another, but the determination of critical values must be made in the context of linkage disequilibrium among SNPs. As there are no reports of allelic variations in IL-1 gene cluster from our subcontinent, the present study was performed to assess interleukin-1 (IL-1) genotype frequencies in our north Indian population, focusing on the IL-1 $\beta$  (promoter&exon-5) and IL-1Ra genes.

## Materials and Methods

### Subjects


The study involved 206 subjects (males 110 & females 96; age range 22-58 yrs) from the north Indian population. Unrelated healthy subjects for the study were drawn from the general population from the same geographical region. The hospital ethical committee approved the study and informed consent was obtained from the participating volunteers.

### DNA Extraction

Five ml of blood was collected in EDTA vials and DNA was extracted from blood lymphocytes using a 'salting out' method (Miller et al., 1988).

### Genotyping of IL-1 Gene Cluster

All subjects were genotyped for three polymorphisms in the interleukin-1 gene cluster: IL-1 $\beta$  promoter region -511, IL-1 $\beta$  exon 5 and IL-1Ra in intron 2, the details of which are presented in Table 1. The size of PCR product was determined using a 100-bp DNA ladder (Roche, Germany). The molecular weight of each band was determined by using software in Alpha Imager 1220 version 5.5 programme where the unknown samples were compared with the 100 bp DNA ladder.



**Figure 1. Schematic picture of the IL-1 gene map demonstrating the different restriction sites for IL-1 $\beta$  promoter region (-511), exon-5 (+3953) restriction enzymes and VNTR in IL-1Ra**

### Statistical Analysis

Two-tailed Fisher's exact test and chi square test was done to compare the allelic frequencies of different populations using the computer software SPSS for windows (version 11.5). P value <0.05 were considered statistically significant. SNPAnalyzer (web-based software) was used to examine Hardy-Weinberg equilibrium (HWE), haplotype estimation by maximum likelihood method, using expectation maximization algorithm and pair wise linkage disequilibrium between each pair of IL-1 loci. (Yoo et al., 2005).

**Table 1. Main Characteristics of Cytokine Gene Polymorphisms and Techniques Used for Screening**

	IL-1 $\beta$ (promoter region)	IL-1 $\beta$ (exon-5)	IL-1Ra
Type of polymorphism	Single base C/T	Single base C/T	86-bp VNTR
Site of polymorphism	-511	+3954	Intron-2
PCR primers			
Upstream	5'-TGGCATTGATC-TGGTTCATC-3'	5'-GTTGTCATCAG-ACTTTGACC-3'	5'-CTCAGCAAC-ACTCCTAT-3'
Downstream	5'-GTTTTAGGAAT-CTTCCACTT-3'	5'-TTCAGTTCAT-ATGGACCAGA-3'	5'-TCCTGGTCTG-CAGGTAA-3'
PCR conditions			
Denaturation	95°C, 0.5 min	95°C, 0.5 min	95°C, 0.5 min
Annealing	56°C, 0.5 min	55°C, 0.5 min	58°C, 0.5 min
Extension	72°C, 0.5 min	72°C, 0.5 min	72°C, 0.5 min
No. of cycles	35	35	30
Digestion	(Ava I)	(Taq I)	86-bp VNTR
Allele size, bp	*C: 190+114 *T: 304	*E1: 135+114 *E2: 249	I: 410, II: 240, III: 500, IV: 325, V: 595, VI: 154

**Table 2. Genotypes and Allele Frequency Distribution of IL-1β (promoter & exon-5) and IL-1Ra Gene Polymorphism in North India**

		Genotypes (%)			Allelic (%), N=412				
IL-1β promoter region									
		CC	CT	TT	C	T			
N = 206		50 (24)	126 (61)	30 (15)	226 (55)	186 (45)			
IL-1β exon-5									
		E1E1	E1E2	E2E2	E1	E2			
N = 206		149(72)	46 (22)	11(6)	344 (84)	68 (16)			
Genotypes (%)									
IL-1Ra									
	1/1(410)	2/2(240)	1/2(410/240)	3/3(500/500)	4/4(325/325)	1/3(410/500)	1/4(410/325)	2/4 (240/325)	
N=206	107(52)	36(18)	50(24)	4(2)	1(0.5)	6(3)	1(0.5)	1(0.5)	
Allele frequency (%)									
	Allele-1	Allele-2	Allele-3	Allele-4					
N= 412	271(66)	123 (30)	14(3)	4 (1)					

**Results**

A significant p-value (p<0.001) revealed that north Indian population is in HWE at all three polymorphic sites of the IL-1 gene i.e. the IL-1β (promoter & exon-5) and IL-1Ra. A schematic IL-1 gene map showing the different restriction sites for IL-1β promoter region (-511), exon-5 (+3953) restriction enzymes and VNTR in IL-1Ra is demonstrated in Fig. 1. Genotype and allele frequencies of IL-1β gene (promoter & exon-5) and IL-1Ra in the subjects are presented in Table 2 and 3 respectively. The observations made for allelic frequency were as follows: (55% vs 45%), (84% vs 16%) and (66%, 30%, 3% & 1%) for (C vs T) for IL-1β promoter, (E1 vs E2) for exon-5 and (1, 2, 3 & 4) for IL-1Ra allele and the percentage of genotypes CC, CT and TT as

24%, 61% and 15%, E1E1, E1E2 and E2E2 as 72%, 22% and 6% and for 1/1, 1/2, 2/2 and ‘others’ (the rare genotypes like 4/4, 1/4, 1/3 & 2/4 were combined as ‘others’) as 52%, 18%, 24% and 6% respectively. The frequency distribution of different genotypes and alleles of IL-1 gene with different populations with reference to ours were compared (Table 4, 5 & 6) by using χ<sup>2</sup> tests. In case of IL-1β -511 (promoter region) significant difference was observed between Portugal, UK, Netherlands, Turkey, Caucasians, African-American, Austria and Egypt as compared to our population. Significant distribution of different genotypes frequency as reported earlier by several studies was observed in IL-1β +3953 (exon-5) polymorphism in Germany, China, Finland, Greece, Japan, Taiwan, Netherlands, Turkey and Egypt population. IL-1Ra, genotype distribution too, was

**Table 3 Genotypes and Allele Frequency Distribution of IL-1β -511 (promoter region) Gene Polymorphism in Various Populations and p-values in Comparison to North Indian Population**

Country/ Ethnicity	No	Age (years)	Genotype (%)			P	Allele (%)			Reference
			(CC)	(CT)	(TT)		(C)	(T)	P	
North India	206	22-58	24	61	15	Ref	55	45	Ref	Present study
Portugal	218	19-61	46	40	14	**	66	34	NS	Machado et al, 2001
UK	54	23-85	41	48	11	*	65	35	NS	Carter et al, 2004
Germany	228	-	39	49	12	NS	64	36	NS	Grimm et al, 2004
Italy	272	59-76	36	44	20	NS	58	42	NS	Zienolddiny et al, 2004
China	361	21.2±1.4	34	48	18	NS	58	42	NS	Zeng et al, 2003
Finland	400	18-60	36	46	18	NS	59	41	NS	Makela et al, 2001
Taiwan	103	50-83	29	47	24	NS	53	47	NS	Tsai et al, 2004
United State (Whites)	124	-	35	50	15	NS	66	34	NS	Cantagrel et al, 1999
Korea	126	58.3±9.6	26	56	18	NS	54	46	NS	Lee et al, 2004
Netherlands	256	-	58	36	6	***	76	24	**	Stokkers et al, 1998
Spain	81	-	32	49	19	NS	57	43	NS	Pastor et al, 2005
Japan	160	40-82	27	49	24	NS	52	48	NS	Nishimura et al, 2002
Turkey	163	35.8±8.4	41	32	27	***	57	43	NS	Coskun et al, 2005
Caucasian	99	-	46	42	12	**	67	33	NS	Wetmore et al, 2005
African-American	98	-	11	58	31	**	40	60	*	Wetmore et al, 2005
Austria	310	37.3±14	42	47	11	*	66	34	NS	Westphal et al, 2003
Egypt	72	59±1.0	37	41	22	*	58	42	NS	Hegab et al, 2005

\* = p<0.05, \*\* = p<0.01, \*\*\* = p<0.001 , at 5 % level of significance

NS = Not Significant (p>0.05)

**Table 4 Genotypes and Allele Frequency Distribution of the IL-1β +3953 (exon-5) Polymorphism and p-values for Different Populations in Comparison to the North Indian Population**

Country/ Ethnicity	No	Age (Years)	Genotype (%)				P	Allele (%)			Reference
			E1E1	E1E2	E2E2	P		E1	E2	P	
North India	206	22-58	72	22	6	Ref	84	16	Ref	Present study	
UK	54	23-85	61	35	4	NS	79	21	NS	Carter et al, 2004	
Germany	228	-	52	44	4	*	74	26	NS	Grimm et al, 2004	
China	361	21.2±1.4	88	12	0	**	94	6	*	Zeng et al, 2003	
Finland	400	-	52	41	7	*	72	28	NS	Maury et al, 2004	
Caucasian	99	-	66	33	1	NS	83	17	NS	Wetmore et al, 2005	
African-American	98	-	78	20	2	NS	88	12	NS	Wetmore et al, 2005	
Greece	110	20-35	50	42	8	**	71	29	*	Sakellari et al, 2003	
Austria	310	37.3±14	59	35	6	NS	77	23	NS	Westphal et al, 2003	
Spain	81	-	61	37	2	NS	79	21	NS	Pastor et al, 2005	
Japan	160	40-82	93	7	0	***	97	3	**	Nishimura et al, 2002	
Taiwan	116	16-28	97	3	0	***	99	1	***	Hang et al, 2003	
Netherlands	131	55-65	55	37	8	*	74	26	NS	Meulenbelt et al, 2004	
Turkey	163	35.8±8	39	42	19	***	60	40	***	Coskun et al, 2005	
Egypt	72	59±1.0	52	41	7	**	73	27	NS	Hegab et al, 2005	

\* = p<0.05, \*\* = p<0.01, \*\*\* = p<0.001 , at 5% level of significance

NS = Not Significant (p>0.05)

significantly different in Portugal, UK, Germany, Japan, Finland, China, Taiwan, Netherlands, Mexico, Korea, Austria and Egypt. We also examined the patterns of LD and haplotype distribution in our population, combining the three sites of IL-1 candidate genes using SNPAnalyzer. All the eight possible haplotypes were common in normals whose frequencies were higher than 1%. Haplotype ‘T-E1-1’ was 41.7% while the least common 2% was observed for ‘C-E2-2’. Linkage disequilibrium was found to be significant between IL-1Ra and IL-1β (promoter region and exon-5) polymorphism in the controls ( $D' = 0.42, p < 0.001$ ).

**Discussion**

Inheritance of polymorphic cytokine gene alleles is dramatically influenced by ethnicity (Hoffmann et al., 2002). Polymorphisms in cytokine genes may result in inter-individual variation in transcriptional regulation, and thus

in differential cytokine production. It has been widely hypothesized that genetic variants of cytokines could have phenotypic relevance and influence an individual’s internal microenvironment (Loughrey et al., 1998, Saurez et al., 2003). In the present study, we investigated IL-1 gene cluster (IL-1β promoter, exon-5 and IL-1Ra) polymorphisms in the north Indian population and compared with the genotypes reported in different populations worldwide. The variation in our Indian population from the rest of the world population signifies the impact of ethnicity. In our population all the eight possible haplotypes were common and their frequency was > 1%, haplotype ‘T-E1-1’ being the most common (41%) The three polymorphic sites of IL-1 gene cluster were in strong linkage disequilibrium, which suggested that the three variants (IL-1β promoter, exon-5 and IL-1Ra) are strongly associated. .

Epidemiologic studies continue to suggest the significant contribution of genetic variation to cancer susceptibility.

**Table 5. Genotypes and Allele Frequency Distribution of IL-1Ra Polymorphism and p-values for Different Populations in Comparison to the North Indian Population**

Country/ Ethnicity	No	Age (Years)	Genotype (%)					P	Allele (%)				Reference
			1/1	1/2	2/2	others	1		2	O	P		
North India	206	22-58	52	18	24	6	Ref	66	30	4	Ref	Present study	
Portugal	220	19-61	51	37	9	3	**	70	27	3	NS	Machado et al, 2001	
UK	54	23-85	61	32	7	0	**	77	23	0	NS	Carter et al, 2004	
Germany	234	-	46	41	6	7	***	69	27	4	NS	Hacker et al, 1997	
Japan	160	40-82	91	6	0	3	***	94	3	3	***	Nishimura et al, 2002	
Finland	400	18-60	50	39	9	2	**	70	28	2	NS	Makela et al, 2001	
China	361	21.2±1.4	93	4	1	2	***	95	3	2	***	Zeng et al, 2003	
Taiwan	103	50-83	86	10	1	3	***	91	6	3	***	Tsai et al, 2004	
Netherland	130	55-65	55	37	8	0	**	74	26	0	NS	Meulenbelt et al, 2004	
Mexico	215	18-92	41	44	12	3	**	63	34	3	NS	Garza et al, 2005	
Korea	434	21-72	90	7	0	3	***	94	3	3	***	Chang et al, 2005	
Austria	310	37.3±14	51	35	10	4	**	69	27	4	NS	Westphal et al, 2003	
Egypt	72	59±1	69	21	10	0	**	80	20	0	*	Hegab et al, 2005	

\* = p<0.05, \*\* = p<0.01, \*\*\* = p<0.001 , at 5% level of significance

NS = Not Significant (p>0.05)

**Table 6. Association Analysis of Haplotypes in Three LD Blocks (IL-1 $\beta$  promoter, exon-5 and IL-1Ra genes) between North Indian ESRD Patients and Controls**

No.	Haplotype	Control (n=206)	
		N	frequency
1.	T-E1-1	86	0.4170
2.	C-E1-1	42	0.2058
3.	T-E1-2	31	0.1454
4.	C-E2-1	19	0.0945
5.	C-E1-2	11	0.0516
6.	T-E2-1	8	0.0414
7.	T-E2-2	5	0.0241
8.	C-E2-2	5	0.0203

(C, T, E1, E2 and 1,2 represent the alleles of IL-1 $\beta$  promoter, exon-5 and IL-1Ra genes)

Although involvement of numerous genetic factors in the pathogenesis of neoplasia is well proven, understanding of the complex molecular mechanisms underlying neoplastic growth is still disappointingly incomplete (Alexandre et al., 2004). Most human cancers are characterized by genomic instability, which is the result of the accumulation of multiple genetic alterations and allelic imbalance throughout the genome (Quek et al., 2004). A wide array of studies has further demonstrated differences in cytokine gene polymorphisms depending on ethnicity (Scarel-Caminaga et al., 2002 and Uboldi et al., 2003). Recently, association studies in humans have shown that polymorphisms of the IL-1 gene were associated with several cancers (Kato et al., 2001 and Ito et al., 2002). Our recently published study reported the association of IL-Ra gene polymorphism with prostate cancer (Mittal et al., 2004).

Thus this kind of study may form the basis for future establishment of epidemiological and clinical databases. However, we are in the initial stages of characterising the tools (i.e., the single-nucleotide polymorphism, SNP) in rigorous analysis of the genetic contributions to complex diseases, such as cancer. Individual SNPs may serve as signposts for disease genes and haplotypes are believed to be superior for this purpose. Moreover, linkage analysis is comprehensive and locates genes that exert a major effect on disease susceptibility. This type of research has the initiative of developing a haplotype map of the human genome, the purpose of which is to relate human genetic variation with disease predisposition, specifically cancer. Thus, it is possible that differences in interleukin-1 distribution between north Indian healthy population and other ethnic groups may reflect a unique profile that may affect disease predisposition and prevalence.

## Acknowledgement

The authors are grateful to the Director and Head of the Department, Urology, Sanjay Gandhi Institute of Medical Sciences for providing the necessary facilities. Ms. P.K. Manchanda thanks the Council of Scientific and Industrial Research New Delhi for awarding a junior research fellowship.

## References

- Alexandre Loktionov (2004). Common gene polymorphisms, cancer progression and prognosis. *Cancer Letters*, **208**, 1-33.
- Anderson DW (2001). Cytokines as drug targets. *Drugs*, **4**, 375-7.
- Bioque G, Crusius JBA, Koutroubakis I, et al (1995). Allelic polymorphism in IL-1 $\beta$  and IL-1 receptor antagonist (IL-1Ra) genes in inflammatory bowel disease. *Clin Exp Immunol*, **102**, 379-83.
- Cantagrel A, Navaux F, Loubet-Lescoulie P, et al (1999). Interleukin-1 beta, interleukin-1 receptor antagonist, interleukin-4 and interleukin-10 gene polymorphisms: relationship to occurrence and severity of rheumatoid arthritis. *Arthritis Rheum*, **42**, 1093-100.
- Carter MJ, Jones S, Camp NJ, et al (2004). Functional correlates of the interleukin-1 receptor antagonist gene polymorphism in the colonic mucosa in ulcerative colitis. *Genes Immun*, **5**, 8-15.
- Chang YW, Jang JY, Kim NH, et al (2005). Interleukin-1B (IL-1B) polymorphisms and gastric mucosal levels of IL-1beta cytokine in Korean patients with gastric cancer. *Int J Cancer*, **114**, 465-71.
- Coskun M, Bacanlı A, Sallakci N, et al (2005). Specific interleukin-1 gene polymorphisms in Turkish patients with Behcet's disease. *Exp Dermatol*, **14**, 124-9.
- Garza-Gonzalez E, Bosques-Padilla FJ, El-Omar E, et al (2005). Role of the polymorphic IL-1B, IL-1RN and TNF-A genes in distal gastric cancer in Mexico. *Int J Cancer*, **114**, 237-41.
- Grimm C, Berger I, Tomovski C, et al (2004). A polymorphism of the interleukin-1 receptor antagonist plays a prominent role within the interleukin-1 gene cluster in vulvar carcinogenesis. *Gynecol Oncol*, **92**, 936-40.
- Hacker UT, Gomolka M, Keller E, et al (1997). Lack of association between an interleukin-1 receptor antagonist gene polymorphism and ulcerative colitis. *Gut*, **40**, 623-7.
- Hegab AE, Sakamoto T, Saitoh W, et al (2005). Polymorphisms of TNFalpha, IL1beta, and IL1RN genes in chronic obstructive pulmonary disease. *Biochem Biophys Res Commun*, **329**, 1246-52.
- Hoffmann SC, Stanley EM, Cox ED, et al (2002). Ethnicity greatly influences cytokine gene polymorphism distribution. *Am J Transpl*, **2**, 560-7.
- Hu Z, Shao M, Chen Y, et al (2005). Allele 2 of the interleukin-1 receptor antagonist gene (IL1RN\*2) is associated with a decreased risk of primary lung cancer. *Cancer Lett*, [Epub ahead of print].
- Ito LS, Iwata H, Hamajima N, et al (2002). Significant reduction in breast cancer risk for Japanese women with interleukin 1 B-31 CT/TT relative to CC genotype. *Jpn J Clin Oncol*, **32**, 398-402.
- Kato S, Onda M, Yamada S, et al (2001). Association of the interleukin-1 beta genetic polymorphism and gastric cancer risk in Japan. *J Gastroenterol*, **36**, 696-99.
- Konishi N, Miki C, Yoshida T, et al (2005). Interleukin-1 receptor antagonist inhibits the expression of vascular endothelial growth factor in colorectal carcinoma. *Oncology*, **68**, 138-45.
- Lee SH, Ihm CG, Sohn SD, et al (2004). Polymorphisms in interleukin-1 beta and Interleukin-1 receptor antagonist genes are associated with kidney failure in Korean patients with type 2 diabetes mellitus. *Am J Nephrol*, **24**, 410-4.
- Loughrey BV, Maxwell AP, Fogarty DG, et al (1998). An interleukin 1B allele, which correlates with a high secretor phenotype, is associated with diabetic nephropathy. *Cytokine*, **10**, 984-8.

- Machado JC, Pharoah P, Sousa S, et al (2001). Interleukin 1B and interleukin 1RN polymorphisms are associated with increased risk of gastric carcinoma. *Gastroenterology*, **121**, 823-9.
- Makela S, Hurme M, Ala-Houhala I, et al (2001). Polymorphism of the cytokine genes in hospitalized patients with Puumala hantavirus infection. *Nephrol Dial Transplant*, **16**, 1368-73.
- Maury CP, Liljestrom M, Laiho K, et al (2004). Anaemia of chronic disease in AA amyloidosis is associated with allele 2 of the interleukin-1beta-511 promoter gene and raised levels of interleukin-1beta and interleukin-18. *J Intern Med*, **256**, 145-52.
- Michael C Iannuzzi, Mary Maliarik, Benjamin Rybicki (2002). Genetic polymorphisms in lung disease: bandwagon or breakthrough? *Respiratory Research*, **3**, 15-22.
- Miller SA, Dykes DD, Polesky HF (1988). A simple salting out procedure for extracting DNA from human nucleated cells. *Nucleic Acids Res*, **16**, 1215.
- Mittal RD, Mishra DK, Bid HK, et al (2004). Interleukin-1 Receptor Antagonist Polymorphism in Patients with Prostate Cancer and Benign Prostatic Hyperplasia: A Case Control Study from North India. *UroOncology*, **4**, 131-4.
- Nishimura M, Kawakami H, Komure O, et al (2002). Contribution of the interleukin-1beta gene polymorphism in multiple system atrophy. *Mov Disord*, **17**, 808-11.
- Pastor IJ, Laso FJ, Romero A, et al (2005). Interleukin-1 gene cluster polymorphisms and alcoholism in Spanish men. *Alcohol Alcohol*, **40**, 181-6.
- Quek ML, Sanderson K, Daneshmand S, et al (2004). New molecular markers for bladder cancer detection. *Curr Opin Urol*, **14**, 259-64.
- Scarel-Caminaga RM, Trevilatto PC, Souza AP, et al (2002). Frequencies of the -330 (T → G) IL-2 and -590 (T → C) IL-4 gene polymorphisms in a population from south-eastern Brazil. *Eur J Immunogenet*, **29**, 293-6.
- Stokkers PC, van Aken BE, Basoski N, et al (1998). Five genetic markers in the interleukin 1 family in relation to inflammatory bowel disease. *Gut*, **43**, 33-9.
- Suarez A, Castro P, Alonso R, et al (2003). Interindividual variations in constitutive interleukin-10 messenger RNA and protein levels and their association with genetic polymorphisms. *Transplantation*, **75**, 711-7.
- Tsai Y-Y, Lee H, Tseng S-H, et al (2004). Evaluation of TNF- $\alpha$  and IL-1 $\beta$  polymorphisms in Taiwan Chinese patients with pterygium. *Eye*, 1-4.
- Uboldi de Capei MU, Dametto E, Fasano ME, et al (2003). Genotyping for cytokine polymorphisms: allele frequencies in the Italian population. *Eur J Immunogenet*, **30**, 5-10.
- Westphal GA, Schnuch A, Moessner R, et al (2003). Cytokine gene polymorphisms in allergic contact dermatitis. *Contact Dermatitis*, **48**, 93-8.
- Wetmore JB, Hung AM, Lovett DH, et al (2005). Interleukin-1 gene cluster polymorphisms predict risk of ESRD. *Kidney Int*, **68**, 278-84.
- Yoo J, Seo B, Kim Y (2005). SNPAnalyzer: a web-based integrated workbench for single-nucleotide polymorphism analysis. *Nucleic Acids Research*, **33**, Web Server issue W483-W488.
- Zeng ZR, Hu PJ, Hu S, et al (2003). Association of interleukin 1B gene polymorphism and gastric cancers in high and low prevalence regions in China. *Gut*, **52**, 1684-9.
- Zienolddiny S, Ryberg D, Maggini V, Skaug V, et al (2004). Polymorphisms of the interleukin-1 beta gene are associated with increased risk of non-small cell lung cancer. *Int J Cancer*, **109**, 353-6.